

# Irradiation induces bone injury by damaging bone marrow microenvironment for stem cells

Xu Cao<sup>a,1</sup>, Xiangwei Wu<sup>a,b</sup>, Deborah Frassica<sup>c</sup>, Bing Yu<sup>a</sup>, Lijuan Pang<sup>a,b</sup>, Lingling Xian<sup>a</sup>, Mei Wan<sup>a</sup>, Weiqi Lei<sup>a</sup>, Michael Armour<sup>d</sup>, Erik Tryggestad<sup>d</sup>, John Wong<sup>d</sup>, Chun Yi Wen<sup>e</sup>, William Weijia Lu<sup>e</sup>, and Frank J. Frassica<sup>a,1</sup>

Departments of <sup>a</sup>Orthopaedic Surgery, <sup>c</sup>Radiation Oncology and Molecular Radiation Sciences, and <sup>d</sup>Radiation Oncology Medical Physics, Johns Hopkins University School of Medicine, Baltimore, MD 21205; <sup>b</sup>Shihezi Medical College, Shihezi University, Shihezi Xinjiang 832002, China; and <sup>e</sup>Department of Orthopaedics, University of Hong Kong, Hong Kong 999777, China

Edited\* by Peter N. Devreotes, Johns Hopkins University School of Medicine, Baltimore, MD, and approved December 16, 2010 (received for review October 14, 2010)

Radiation therapy can result in bone injury with the development of fractures and often can lead to delayed and nonunion of bone. There is no prevention or treatment for irradiation-induced bone injury. We irradiated the distal half of the mouse left femur to study the mechanism of irradiation-induced bone injury and found that no mesenchymal stem cells (MSCs) were detected in irradiated distal femora or nonirradiated proximal femora. The MSCs in the circulation doubled at 1 week and increased fourfold after 4 wk of irradiation. The number of MSCs in the proximal femur quickly recovered, but no recovery was observed in the distal femur. The levels of free radicals were increased threefold at 1 wk and remained at this high level for 4 wk in distal femora, whereas the levels were increased at 1 wk and returned to the basal level at 4 wk in nonirradiated proximal femur. Free radicals diffuse ipsilaterally to the proximal femur through bone medullary canal. The blood vessels in the distal femora were destroyed in angiographic images, but not in the proximal femora. The osteoclasts and osteoblasts were decreased in the distal femora after irradiation, but no changes were observed in the proximal femora. The total bone volumes were not affected in proximal and distal femora. Our data indicate that irradiation produces free radicals that adversely affect the survival of MSCs in both distal and proximal femora. Irradiation injury to the vasculatures and the microenvironment affect the niches for stem cells during the recovery period.

CFU-fibroblast | repopulation | differentiation | self-renewal | angiography

Ionizing radiation is often used to treat malignancies such as breast cancer, lymphoma, Ewing sarcoma, soft tissue sarcomas, and rectal and anorectal carcinomas. Bone insufficiency is a significant problem resulting in stress fractures, bone fragmentation, and joint dissolution (1, 2). The risk of a hip fracture for a woman following pelvic irradiation for the treatment of carcinomas of the anus, cervix, or rectum is as much as three times that of the population of women who do not receive irradiation (3). Women older than the age of 65 y have a 16% lifetime risk of a hip fracture. Fractures of the sacrum are also very common following irradiation. These stress or insufficiency fractures are difficult to treat, with very high rates of delayed union and nonunion of the fractures. Only small amounts of bone healing (i.e., callus) is noted on serial radiographs. Surgical treatment with internal fixation and conventional bone grafting has had only limited success. The mechanism of irradiation-induced bone injury has not been fully elucidated.

Human bone marrow contains both hematopoietic stem cells (HSCs) and the stromal cell compartments; the latter comprises various cell types, including mesenchymal stem cells (MSCs) (4, 5). The MSCs derived from adult bone marrow are pluripotent and can differentiate into heart, lung, and gut epithelia in the animal (6–8). Repopulation with bone marrow-derived MSCs could lead to differentiation into different cell types, including osteoblasts, adipocytes, chondrocytes, and fibroblasts (5). The microenvironment in the bone marrow comprises vascular niche

and stromal cell niche including osteoblasts for stem cell self-renewal and differentiation (9, 10). Bone marrow is extremely vulnerable to cytotoxicity caused by radiation therapy (11). In contrast to HSCs, very few data have been published on the radiosensitivity of MSCs. Several recent studies have reported that MSCs from patients exhibit a complete host profile after total body irradiation and bone marrow transplantation, whereas cells differentiated from HSCs are of donor origin (12, 13). However, the cellular and molecular mechanisms accounting for such radiation tolerance in MSCs remain elusive.

Radiation generates free radicals and reactive oxygen species, which damage vital cellular targets such as DNA and membranes, and affect the niches for stem cells (14). Both HSCs and MSCs are implicated in the repair of irradiation damage of distal epithelial through their mobilization in the circulation (15–17). In this study, we aimed to investigate how irradiation affects bone marrow stem cells and residing microenvironment directly or indirectly by using a unique Small Animal Radiation Research Platform (SARRP) built by the Department of Radiation Oncology at Johns Hopkins University, in which computed tomography (CT), X-ray, and irradiation are integrated into a single system (18). We found that irradiation damaged the bone marrow microenvironment for stem cells, and our findings suggest that prevention of the damage during radiation therapy will be helpful for the recovery of bone injury.

## Results

To investigate the mechanism of irradiation-induced bone injury, we irradiated mouse femora by using a unique SARRP (Fig. 1A). The irradiation can be delivered at an accuracy of 1 mm resolution with the aid of micro-CT and X-ray. Groups of mice were irradiated at the distal half of the femora with a dose of 4 Gy once per day for 5 d consecutively. The left half of the proximal femora and the right femora did not receive irradiation as controls (Fig. 1B and C). Histologic staining of the femoral sections showed that irradiation induced adipogenesis in the irradiated left distal femur at 1 wk relative to the right nonirradiated femur. Adipocytes were significantly increased at 4 wk after radiation in the medullary cavity (Fig. 1D). The results indicate that irradiation changes the bone marrow microenvironment.


We then examined whether irradiation injured bone and bone cells directly. The total bone volumes were not affected in both the left proximal and distal femora 1 or 4 wk after irradiation in CT analysis (Fig. 2A), and the trabecular bone thickness, num-

Author contributions: X.C., X.W., and F.J.F. designed research; D.F., B.Y., L.P., L.X., M.W., W.L., M.A., E.T., and C.Y.W. performed research; J.W. and W.W.L. contributed new reagents/analytic tools; X.C., X.W., and F.J.F. analyzed data; and X.C., X.W., and F.J.F. wrote the paper.

The authors declare no conflict of interest.

\*This Direct Submission article had a prearranged editor.

<sup>1</sup>To whom correspondence may be addressed. E-mail: ffrassi@jhmi.edu or xcao11@jhmi.edu.



**Fig. 1.** Local Irradiation on left distal femur. (A) SARRP system. (B) Radiograph of left lower limb of a mouse that received local irradiation of the distal femur. Light rectangle indicates the beam of X-ray. (C) Femora collected from the mice 1 wk after irradiation. (D) Light micrographs of H&E staining performed on trabecular bone sections from proximal and distal femora of mice 1 wk and 4 wk after irradiation. Yellow cross indicates left proximal femur, green cross indicates irradiated area of left distal femur, yellow triangle indicates right proximal femur, and green triangle indicates right distal femur. L, left, R, right. (Scale bar: 100  $\mu\text{m}$ .)


ber, and separation were not changed either (Fig. 2 *B–D*). Bone histomorphometry analysis showed that numbers of both osteoclasts and osteoblasts were decreased in the distal femora after irradiation, but no changes were observed in the proximal femora (Fig. 2 *E* and *F*). The results indicate that irradiation affects osteoblasts and osteoclasts in the distal femora, but has no significant effect on differentiated osteoclasts and osteoblasts in the proximal femora (i.e., nonirradiated area).

The effects of irradiation on bone marrow MSCs were examined in CFU-fibroblast (CFU-F) assay with bone marrow cells isolated from left distal femora (i.e., irradiated area) or left proximal femora (i.e., nonirradiated area). No colonies of MSCs were observed in the left proximal femur or distal femur at 1 wk after irradiation, whereas colonies of CFU-F and CFU-osteoblast (CFU-Ob) were observed in the nonirradiated right femur (Fig. 3 *A* and *B*). Interestingly, recovery of MSCs was observed in the left proximal femur at 4 wk, but not for irradiated distal femur. Similar results were observed and osteoblast differentiation potential of MSCs in CFU-Ob assay (Fig. 3 *A* and *C*). The colonies of MSCs in the left and right nonirradiated tibia were not changed (Fig. 3 *D*). MSCs have been shown to be mobilized in the circulation in response to tissue damage. We therefore examined the effects of radiation on the levels of MSCs in the circulation. Previously, we have demonstrated the  $\text{CD}29^+\text{Sca-1}^+\text{CD}45^-\text{CD}11\text{b}^-$  bone marrow stroma are subset of MSCs (19). Flow cytometry analysis of  $\text{CD}29^+\text{Sca-1}^+\text{CD}45^-\text{CD}11\text{b}^-$  MSCs in peripheral blood collected from the mice at 1 and 4 wk after treatment in the irradiation and sham groups indicates that  $\text{CD}29^+\text{Sca-1}^+\text{CD}45^-\text{CD}11\text{b}^-$  MSCs were doubled at 1 wk after irradiation and increased fourfold in 4 wk (Fig. 3 *E*). The level of  $\text{CD}29^+\text{Sca-1}^+\text{CD}45^-\text{CD}11\text{b}^-$  MSCs was also increased in proxi-

mal femur at 4 wk after irradiation whereas the levels in the right femur and left distal femur were unchanged. The results reveal that bone marrow MSCs can be affected indirectly following irradiation at another site in the bone marrow cavity. The increased level of Sca-1-positive MSCs in response to irradiation suggests that there is a systemic response resulting in mobilization of MSCs for the repair of bone injury.

To examine the mechanism of irradiation-induced damage of bone marrow MSCs, we measured the levels of free radicals in the bone marrow isolated from the irradiated left distal femur, nonirradiated proximal left proximal femur, and right femur by using thiobarbituric acid reactive substances (TBARS) assay. This assay measures the consequences of lipid peroxidation and malondialdehyde specifically. The levels of TBARS were increased threefold at 1 wk and remained at the same level for 4 wk in left distal femora (Fig. 4 *A*), whereas the levels were increased at 1 wk and decreased to the basal level at 4 wk in nonirradiated proximal femur (Fig. 4 *A*). No changes of TBARS were observed in the right femora (Fig. 4 *A*) and circulation (Fig. 4 *B*), indicating that the free radicals diffuse to ipsilaterally distal femur through bone medullary channel, leading to MSCs damage. The results also suggest that irradiation produces free radicals that adversely affect the MSCs in both distal and proximal femora, but have no significant effects on differentiated osteoclasts and osteoblasts in the ipsilateral proximal femur (i.e., nonirradiated area).

Vasculatures are essential for the maintenance of MSCs as their niche (20, 21). Bone formation is coupled with angiogenesis (22–24). Angiographic images of microphil-perfused femora were performed to analyze the effects of irradiation on bone marrow vasculatures. The results showed that the blood vessels



**Fig. 2.** Bone histomorphometric analysis of trabecular bone from the mice 1 wk and 4 wk after irradiation. Bone volume fraction (BV/TV) (A), trabecular thickness (Tb.Th) (B), trabecular number (Tb.N) (C), trabecular separation (Tb.Sp) (D), number of osteoblasts per bone perimeter (N.Ob\*/B.Pm) (E), and number of osteoclasts per bone perimeter (N.Oc\*/B.Pm) (F) were measured (LP, left proximal femur; LD, left distal femur; RP, right proximal femur; RD, right distal femur). Data represent the mean  $\pm$  SEM;  $n = 10$  (\* $P < 0.05$ ).

in the irradiated left distal femora were damaged at 1 wk and 4 wk after irradiation, but the vessels were intact in the non-irradiated left proximal femora (Fig. 4C). Morphometric analysis showed that vessel volume, thickness, and number were decreased in the distal femur relative to controls (Fig. 4D–F). The results suggest that direct irradiation, not free radical generation, damages the bone marrow vasculatures.


### Discussion

In this study, precise irradiation of the mouse distal femur was performed with a unique SARRP built with integration of CT, X-ray, and irradiation in a single treatment platform. The mouse distal half of the femur was precisely irradiated at accuracy of 1 mm. Here, the direct and indirect effects of irradiation on bone injury and bone marrow MSCs were examined in the same mouse femur with common bone marrow cavity. Our results reveal that irradiation can affect bone marrow MSCs indirectly in the proximal femur without direct irradiation. Interestingly, no colony formation of MSCs was observed in CFU-F and CFU-Ob assays in irradiated and nonirradiated areas of the bone marrow, and the levels of free radicals were dramatically increased. Most likely, the free radicals produced during radiation affects the survival, self-renewal, or differentiation of MSCs. Interestingly, the recovery of MSCs colony formation was seen in 4 wk after irradiation in the nonirradiated proximal femur, but not in the irradiated distal femur. Importantly, the vasculatures—the potential niche for stem cells—were destroyed in irradiated areas. Damage of bone marrow vasculatures—the niches for stem cells—likely affects the recovery of MSCs in the irradiated areas, which may explain the refractory nature of bone injury and fractures in patients receiving ionizing radiation to bone. Therefore, our findings suggest that irradiation induces bone injury by producing free radicals that adversely affect microenvironment for MSCs and damaging bone marrow blood vessels.

Bone injury and fractures often occur in patients receiving curative doses of ionizing radiation. These fractures often heal very slowly and sometimes fail to unite. These clinical observations are in accord with our findings in this mouse model and suggest that irradiation damages the bone marrow microenvironment or the niches for stem cells. Recovery of bone injury from ionizing irradiation may occur from repopulation of inhabitant stem cells (14) or repopulation from the systemic circulation. Our results showed that MSCs in the direct irradiated area did not recover, whereas the MSCs in the nonirradiated proximal left proximal femur recovered quickly. The free radicals produced changed the microenvironment in the bone marrow and likely brought the MSCs into a quiescent phase in the nonirradiated area of bone marrow. Stem cells of hematopoietic and mesenchymal origin have been shown in the recovery of ionizing irradiation-induced distal epithelial damage (15–17). The bone marrow with damaged microvasculature is not likely able to retain stem cells and lead to refractory of bone injury and fractures. It has been implicated that mobilization of stem cells in the circulation can occur in response to irradiation (25–27). Indeed, the levels of Sca-1-positive MSCs were increased in the circulation after irradiation. Different markers have been used for MSCs. A recent study revealed that bone marrow MSCs are nestin-positive cells (28). It will be interesting to investigate whether the mobilized Sca-1-positive MSCs are also nestin-positive and if that contributes to the recovery of MSCs in the bone marrow of proximal femur. It will also be interesting to investigate whether the mobilized Sca-1-positive MSCs contribute to the recovery of MSCs in the bone marrow of proximal femur.

The therapeutic potential of bone marrow-derived MSCs has recently been implicated. Not only are these cells able to migrate to injured tissues, but they also are able to differentiate into different phenotypes according to the tissues in which they reside (29). Furthermore, it has been shown that MSCs of bone marrow





**Fig. 4.** Effects of local irradiation on the production of free radicals and angiogenesis. Free radicals were assessed in both bone marrow (A) and peripheral blood (B) before irradiation and 1 wk and 4 wk after irradiation detected by TBARS assay. (C) Representative 3D micro-CT images of femora from the mice after local irradiation. (Scale bar: 1 mm.). (D–F) Quantitative micro-CT angiography analysis. Vascular volume fraction (VV/TV) (D), vascular thickness (V.Th) (E), and vascular number (V.N) (F) were measured. Data represent the mean  $\pm$  SEM;  $n = 10$  (\* $P < 0.05$ ).

**Micro-CT Angiography Analysis.** Blood vessels in bone were imaged by angiography of microphil-perfused long bones. The thoracic cavity was opened, and the inferior vena cava was severed after animals were killed. The vasculature was flushed with 0.9% normal saline solution containing heparin sodium (100 U/mL) through a needle inserted into the left ventricle. The specimens were pressure fixed with 10% neutral buffered formalin. Formalin was flushed from the vessels by using heparinized saline solution, and the vasculature was injected with a radiopaque silicone rubber compound containing lead chromate (Microfil MV-122; Flow Tech). Samples were stored at 4 °C overnight for contrast agent polymerization. Mouse femurs were dissected from the specimens and soaked for 4 d in 10% neutral buffered

formalin to ensure complete tissue fixation. Specimens were subsequently treated for 48 h in a formic acid-based solution (Cal-Ex II) to decalcify the bone and facilitate image thresholding of the femoral vasculature from the surrounding tissues. Images were obtained using a high resolution (9- $\mu$ m isotropic voxel size) micro-CT imaging system (MicroCT40; Scanco). A threshold of 306 was initially chosen based on visual interpretation of thresholded 2D tomograms.

**Statistics.** Data are presented as mean  $\pm$  SEM and were analyzed using a one-way ANOVA followed by Dunnett test.

- Costantino PD, Friedman CD, Steinberg MJ (1995) Irradiated bone and its management. *Otolaryngol Clin North Am* 28:1021–1038.
- Jegoux F, Malard O, Goyenvallée E, Aguado E, Daculsi G (2010) Radiation effects on bone healing and reconstruction: interpretation of the literature. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 109:173–184.
- Baxter NN, Habermann EB, Tepper JE, Durham SB, Virnig BA (2005) Risk of pelvic fractures in older women following pelvic irradiation. *JAMA* 294:2587–2593.
- Gerson SL (1999) Mesenchymal stem cells: No longer second class marrow citizens. *Nat Med* 5:262–264.
- Prockop DJ (1997) Marrow stromal cells as stem cells for nonhematopoietic tissues. *Science* 276:71–74.
- Jiang Y, et al. (2002) Pluripotency of mesenchymal stem cells derived from adult marrow. *Nature* 418:41–49.
- Kawada H, et al. (2004) Nonhematopoietic mesenchymal stem cells can be mobilized and differentiate into cardiomyocytes after myocardial infarction. *Blood* 104:3581–3587.
- Lee KD, et al. (2004) In vitro hepatic differentiation of human mesenchymal stem cells. *Hepatology* 40:1275–1284.
- Garrett RW, Emerson SG (2009) Bone and blood vessels: The hard and the soft of hematopoietic stem cell niches. *Cell Stem Cell* 4:503–506.
- Yin T, Li L (2006) The stem cell niches in bone. *J Clin Invest* 116:1195–1201.
- Mauch P, et al. (1995) Hematopoietic stem cell compartment: Acute and late effects of radiation therapy and chemotherapy. *Int J Radiat Oncol Biol Phys* 31:1319–1339.
- Dickhut A, et al. (2005) Mesenchymal stem cells obtained after bone marrow transplantation or peripheral blood stem cell transplantation originate from host tissue. *Ann Hematol* 84:722–727.
- Rieger K, et al. (2005) Mesenchymal stem cells remain of host origin even a long time after allogeneic peripheral blood stem cell or bone marrow transplantation. *Exp Hematol* 33:605–611.
- Greenberger JS, Epperly M (2009) Bone marrow-derived stem cells and radiation response. *Semin Radiat Oncol* 19:133–139.
- Epperly MW, et al. (2004) Bone marrow origin of cells with capacity for homing and differentiation to esophageal squamous epithelium. *Radiat Res* 162:233–240.
- Greenberger JS, Epperly MW (2007) Review. Antioxidant gene therapeutic approaches to normal tissue radioprotection and tumor radiosensitization. *In Vivo* 21:141–146.
- Krause DS, et al. (2001) Multi-organ, multi-lineage engraftment by a single bone marrow-derived stem cell. *Cell* 105:369–377.
- Deng H, et al. (2007) The small-animal radiation research platform (SARRP): Dosimetry of a focused lens system. *Phys Med Biol* 52:2729–2740.
- Tang Y, et al. (2009) TGF-beta1-induced migration of bone mesenchymal stem cells couples bone resorption with formation. *Nat Med* 15:757–765.
- Crisan M, et al. (2008) A perivascular origin for mesenchymal stem cells in multiple human organs. *Cell Stem Cell* 3:301–313.
- Shi S, Gronthos S (2003) Perivascular niche of postnatal mesenchymal stem cells in human bone marrow and dental pulp. *J Bone Miner Res* 18:696–704.
- Schipani E, Maes C, Carmeliet G, Semenza GL (2009) Regulation of osteogenesis-angiogenesis coupling by HIFs and VEGF. *J Bone Miner Res* 24:1347–1353.
- Street J, et al. (2002) Vascular endothelial growth factor stimulates bone repair by promoting angiogenesis and bone turnover. *Proc Natl Acad Sci USA* 99:9656–9661.
- Wang Y, et al. (2007) The hypoxia-inducible factor alpha pathway couples angiogenesis to osteogenesis during skeletal development. *J Clin Invest* 117:1616–1626.



## Corrections

### MEDICAL SCIENCES

Correction for “Irradiation induces bone injury by damaging bone marrow microenvironment for stem cells,” by Xu Cao, Xiangwei Wu, Deborah Frassica, Bing Yu, Lijuan Pang, Lingling Xian, Mei Wan, Weiqi Lei, Michael Armour, Erik Tryggestad, John Wong, Chun Yi Wen, William Weijia Lu, and Frank J. Frassica, which appeared in issue 4, January 25, 2011, of *Proc Natl Acad Sci USA* (108:1609–1614; first published January 10, 2011; 10.1073/pnas.1015350108).

The authors note that their conflict of interest statement was omitted during publication. The authors declare that John Wong has a research service contract with Gulmay Medical, Inc. in the transfer of the SARRP technology from Johns Hopkins to the company. The SARRP was used by Dr. Xu Cao to irradiate his study animals, and was only peripherally related to the subject matter of the manuscript. Additionally, the specific unit was constructed at Johns Hopkins with grant support from the National Cancer Institute (US), and not by Gulmay.

[www.pnas.org/cgi/doi/10.1073/pnas.1104040108](http://www.pnas.org/cgi/doi/10.1073/pnas.1104040108)

### DEVELOPMENTAL BIOLOGY

Correction for “Cardiac-specific transcription factor genes *Smad4* and *Gata4* cooperatively regulate cardiac valve development,” by Ivan P. Moskowitz, Jun Wang, Michael A. Peterson, William T. Pu, Alexander C. Mackinnon, Leif Oxburgh, Gerald C. Chu, Molly Sarkar, Charles Berul, Leslie Smoot, Elizabeth J. Robertson, Robert Schwartz, Jonathan G. Seidman, and Christine E. Seidman, which appeared in issue 10, March 8, 2011, of *Proc Natl Acad Sci USA* (108:4006–4011; first published February 17, 2011; 10.1073/pnas.1019025108).

The authors note that the title appeared incorrectly. The title should instead appear as “Transcription factor genes *Smad4* and *Gata4* cooperatively regulate cardiac valve development.” The online version has been corrected.

[www.pnas.org/cgi/doi/10.1073/pnas.1103973108](http://www.pnas.org/cgi/doi/10.1073/pnas.1103973108)